

# The Physics and Cosmology of TeV Blazars

Christoph Pfrommer<sup>1</sup>

in collaboration with

Avery E. Broderick<sup>2</sup>, Phil Chang<sup>3</sup>, Ewald Puchwein<sup>1</sup>, Volker Springel<sup>1</sup>

<sup>1</sup>Heidelberg Institute for Theoretical Studies, Germany

<sup>2</sup>Perimeter Institute/University of Waterloo, Canada


<sup>3</sup>University of Wisconsin-Milwaukee, USA

Jul 11, 2012 / Meeting, DFG Research Unit FOR 1254




# The Hitchhiker's Guide to ... Blazar Heating

- the extragalactic TeV Universe
- plasma physics for cosmologists



# The Hitchhiker's Guide to ... Blazar Heating

- the extragalactic TeV Universe
- plasma physics for cosmologists
- consequences for
  - intergalactic magnetic fields
  - extragalactic gamma-ray background
  - thermal history of the Universe
  - Lyman- $\alpha$  forest
  - “missing dwarf galaxies”
  - H I mass function
  - galaxy cluster bimodality



# Outline

- 1 Physics of blazar heating
  - TeV emission from blazars
  - Plasma instabilities and magnetic fields
  - Extragalactic gamma-ray background
- 2 The intergalactic medium
  - Properties of blazar heating
  - Thermal history of the IGM
  - The Lyman- $\alpha$  forest
- 3 Structure formation
  - Formation of dwarf galaxies
  - Puzzles in galaxy formation
  - Conclusions




# TeV gamma-ray astronomy

H.E.S.S.




MAGIC I



VERITAS



MAGIC II




# The TeV gamma-ray sky

There are several classes of TeV sources:


- Galactic - pulsars, BH binaries, supernova remnants
- Extragalactic - **mostly** blazars, two starburst galaxies

VHE  $\gamma$ -ray Sky Map  
( $E_\gamma > 100$  GeV)




2011-01-08 - Up-to-date plot available at <http://www.mpp.mpg.de/~nwagner/sources/>

# Unified model of active galactic nuclei



# The blazar sequence



Ghisellini (2011), arXiv:1104.0006

- continuous sequence from **L**BL–**I**BL–**H**BL
- TeV blazars are dim (very sub-Eddington)
- TeV blazars have rising spectra in the Fermi band ( $\alpha < 2$ )
- define TeV blazar = **hard I**BL + **H**BL





# Propagation of TeV photons

- 1 TeV photons can pair produce with 1 eV **EBL photons**:

$$\gamma_{\text{TeV}} + \gamma_{\text{eV}} \rightarrow e^+ + e^-$$



# Propagation of TeV photons

- 1 TeV photons can pair produce with 1 eV **EBL photons**:

$$\gamma_{\text{TeV}} + \gamma_{\text{eV}} \rightarrow e^+ + e^-$$

- mean free path for this depends on the density of 1 eV photons:
  - $\lambda_{\gamma\gamma} \sim (35 \dots 700)$  Mpc for  $z = 1 \dots 0$
  - pairs produced with energy of 0.5 TeV ( $\gamma = 10^6$ )



# Propagation of TeV photons

- 1 TeV photons can pair produce with 1 eV **EBL photons**:

$$\gamma_{\text{TeV}} + \gamma_{\text{eV}} \rightarrow e^+ + e^-$$

- mean free path for this depends on the density of 1 eV photons:
  - $\lambda_{\gamma\gamma} \sim (35 \dots 700) \text{ Mpc}$  for  $z = 1 \dots 0$
  - pairs produced with energy of 0.5 TeV ( $\gamma = 10^6$ )
- these pairs inverse Compton scatter off the **CMB photons**:
  - mean free path is  $\lambda_{\text{IC}} \sim \lambda_{\gamma\gamma}/1000$
  - producing gamma-rays of  $\sim 1 \text{ GeV}$

$$E \sim \gamma^2 E_{\text{CMB}} \sim 1 \text{ GeV}$$



# Propagation of TeV photons

- 1 TeV photons can pair produce with 1 eV **EBL photons**:

$$\gamma_{\text{TeV}} + \gamma_{\text{eV}} \rightarrow e^+ + e^-$$

- mean free path for this depends on the density of 1 eV photons:
  - $\rightarrow \lambda_{\gamma\gamma} \sim (35 \dots 700) \text{ Mpc}$  for  $z = 1 \dots 0$
  - $\rightarrow$  pairs produced with energy of 0.5 TeV ( $\gamma = 10^6$ )
- these pairs inverse Compton scatter off the **CMB photons**:
  - $\rightarrow$  mean free path is  $\lambda_{\text{IC}} \sim \lambda_{\gamma\gamma}/1000$
  - $\rightarrow$  producing gamma-rays of  $\sim 1 \text{ GeV}$


$$E \sim \gamma^2 E_{\text{CMB}} \sim 1 \text{ GeV}$$

- each TeV point source should also be a GeV point source



# What about the cascade emission?

Every TeV source should be associated with a 1-100 GeV gamma-ray halo – **not seen!**



# Measuring IGM $B$ -fields from TeV/GeV observations

- TeV beam of  $e^+/e^-$  are deflected out of the line of sight  
reducing the GeV IC flux → **lower limit on  $B$**
- Larmor radius

$$r_L = \frac{E}{eB} \sim 30 \left( \frac{E}{3 \text{ TeV}} \right) \left( \frac{B}{10^{-16} \text{ G}} \right)^{-1} \text{ Mpc}$$



# Measuring IGM $B$ -fields from TeV/GeV observations

- TeV beam of  $e^+/e^-$  are deflected out of the line of sight reducing the GeV IC flux  $\rightarrow$  **lower limit on  $B$**
- Larmor radius

$$r_L = \frac{E}{eB} \sim 30 \left( \frac{E}{3 \text{ TeV}} \right) \left( \frac{B}{10^{-16} \text{ G}} \right)^{-1} \text{ Mpc}$$

- IC mean free path

$$x_{\text{IC}} \sim 0.1 \left( \frac{E}{3 \text{ TeV}} \right)^{-1} \text{ Mpc}$$

- for the associated 10 GeV IC photons the *Fermi* angular resolution is  $0.2^\circ$  or  $\theta \sim 3 \times 10^{-3} \text{ rad}$


$$\frac{x_{\text{IC}}}{r_L} > \theta \rightarrow B \gtrsim 10^{-16} \text{ G}$$



# Missing plasma physics?

How do beams of  $e^+/e^-$  propagate through the IGM?

- plasma processes are important
- interpenetrating beams of charged particles are unstable
- consider the two-stream instability:



- one frequency (timescale) and one length in the problem:

$$\frac{\omega_p}{\gamma} = \sqrt{\frac{4\pi e^2 n_e}{\gamma^2 m_e}}, \quad \lambda_p = \frac{\gamma c}{\omega_p} \sim 10^{14} \text{ cm} \times \left( \frac{\gamma}{10^6} \right) \Big|_{\bar{\rho}(z=0)}$$






# Two-stream instability: mechanism

wave-like perturbation with  $\mathbf{k} \parallel \mathbf{v}_{\text{beam}}$ , longitudinal charge oscillations in background plasma (Langmuir wave):


- initially homogeneous beam- $e^-$ :  
attractive (repulsive) force by potential maxima (minima)
- $e^-$  attain lowest velocity in potential minima  $\rightarrow$  bunching up
- $e^+$  attain lowest velocity in potential maxima  $\rightarrow$  bunching up




# Two-stream instability: mechanism

wave-like perturbation with  $\mathbf{k} \parallel \mathbf{v}_{\text{beam}}$ , longitudinal charge oscillations in background plasma (Langmuir wave):

- beam- $e^+/e^-$  couple in phase with the background perturbation: enhances background potential
- stronger forces on beam- $e^+/e^- \rightarrow$  positive feedback
- exponential wave-growth  $\rightarrow$  instability



# Two-stream instability: energy transfer





- particles with  $v \gtrsim v_{\text{phase}}$ :  
pair energy  $\rightarrow$  plasma waves  $\rightarrow$  growing modes
- particles with  $v \lesssim v_{\text{phase}}$ :  
plasma wave energy  $\rightarrow$  pairs  $\rightarrow$  damped modes



# Oblique instability


$\mathbf{k}$  oblique to  $\mathbf{v}_{\text{beam}}$ : real world perturbations don't choose "easy"  
alignment =  $\sum$  all orientations



Bret (2009), Bret+ (2010)



# Beam physics – growth rates



- consider a light beam penetrating into relatively dense plasma

- maximum growth rate

$$\sim 0.4 \gamma \frac{n_{\text{beam}}}{n_{\text{IGM}}} \omega_p$$

- oblique instability beats IC by two orders of magnitude

Broderick, Chang, C.P. (2012)



# Beam physics – complications . . .

non-linear saturation:

- non-linear evolution of these instabilities at these density contrasts is not known
- expectation from PIC simulations suggest substantial isotropization of the beam
- **assume** that they grow at linear rate up to saturation



# Beam physics – complications . . .

non-linear saturation:

- non-linear evolution of these instabilities at these density contrasts is not known
- expectation from PIC simulations suggest substantial isotropization of the beam
- **assume** that they grow at linear rate up to saturation

→ plasma instabilities dissipate the beam's energy, no (little) energy left over for inverse Compton scattering off the CMB



# TeV emission from blazars – a new paradigm

$$\gamma_{\text{TeV}} + \gamma_{\text{eV}} \rightarrow e^+ + e^- \rightarrow \begin{cases} \text{IC off CMB} & \rightarrow \gamma_{\text{GeV}} \\ \text{plasma instabilities} & \rightarrow \text{heating IGM} \end{cases}$$





# TeV emission from blazars – a new paradigm

$$\gamma_{\text{TeV}} + \gamma_{\text{eV}} \rightarrow e^+ + e^- \rightarrow \begin{cases} \text{IC off CMB} & \rightarrow \gamma_{\text{GeV}} \\ \text{plasma instabilities} & \rightarrow \text{heating IGM} \end{cases}$$

absence of  $\gamma_{\text{GeV}}$ 's has significant implications for ...

- intergalactic  $B$ -field estimates
- $\gamma$ -ray emission from blazars: spectra, background



# TeV emission from blazars – a new paradigm

$$\gamma_{\text{TeV}} + \gamma_{\text{eV}} \rightarrow e^+ + e^- \rightarrow \begin{cases} \text{IC off CMB} & \rightarrow \gamma_{\text{GeV}} \\ \text{plasma instabilities} & \rightarrow \text{heating IGM} \end{cases}$$

absence of  $\gamma_{\text{GeV}}$ 's has significant implications for ...

- intergalactic  $B$ -field estimates
- $\gamma$ -ray emission from blazars: spectra, background


additional IGM heating has significant implications for ...

- thermal history of the IGM: Lyman- $\alpha$  forest
- late time structure formation: dwarfs, galaxy clusters



# Implications for $B$ -field measurements

Fraction of the pair energy lost to inverse-Compton on the CMB:  $f_{\text{IC}} = \Gamma_{\text{IC}} / (\Gamma_{\text{IC}} + \Gamma_{\text{oblique}})$



Broderick, Chang, C.P. (2012)



# Conclusions on $B$ -field constraints from blazar spectra

- it is thought that TeV blazar spectra might constrain IGM  $B$ -fields
- this assumes that cooling mechanism is IC off the CMB + deflection from magnetic fields
- beam instabilities may allow high-energy  $e^+ / e^-$  pairs to self scatter and/or lose energy
- isotropizes the beam – no need for  $B$ -field
- $\lesssim 1\text{--}10\%$  of beam energy to IC CMB photons




# Conclusions on $B$ -field constraints from blazar spectra

- it is thought that TeV blazar spectra might constrain IGM  $B$ -fields
- this assumes that cooling mechanism is IC off the CMB + deflection from magnetic fields
- beam instabilities may allow high-energy  $e^+ / e^-$  pairs to self scatter and/or lose energy
- isotropizes the beam – no need for  $B$ -field
- $\lesssim 1\text{--}10\%$  of beam energy to IC CMB photons

→ **TeV blazar spectra are not suitable to measure IGM  $B$ -fields!**



# TeV blazar luminosity density: today




Broderick, Chang, C.P. (2012)

- collect luminosity of all 23 TeV blazars with good spectral measurements
- account for the selection effects (sky coverage, duty cycle, galactic occultation, TeV flux limit)
- TeV blazar luminosity density is a scaled version ( $\eta_B \sim 0.2\%$ ) of that of quasars!



# Unified TeV blazar-quasar model




Broderick, Chang, C.P. (2012)

Quasars and TeV blazars are:

- regulated by the same mechanism
- contemporaneous elements of a single AGN population: TeV-blazar activity does not lag quasar activity



# Unified TeV blazar-quasar model



Broderick, Chang, C.P. (2012)

Quasars and TeV blazars are:


- regulated by the same mechanism
- contemporaneous elements of a single AGN population: TeV-blazar activity does not lag quasar activity

→ **assume that they trace each other for all redshifts!**






# How many TeV blazars are there?



Hopkins+ (2007)




# How many TeV blazars are there?



Hopkins+ (2007)




# How many TeV blazars are there?



Hopkins+ (2007)



# Fermi number count of “TeV blazars”




Broderick, Chang, C.P. (2012)

- number evolution of TeV blazars that are expected to have been observed by *Fermi* vs. observed evolution
- colors: different flux (luminosity) limits connecting the *Fermi* and the TeV band:

$$L_{\text{TeV},\min}(z) = \eta L_{\text{Fermi},\min}(z)$$

# Fermi number count of “TeV blazars”



Broderick, Chang, C.P. (2012)


- number evolution of TeV blazars that are expected to have been observed by *Fermi* vs. observed evolution
- colors: different flux (luminosity) limits connecting the *Fermi* and the TeV band:

$$L_{\text{TeV},\min}(z) = \eta L_{\text{Fermi},\min}(z)$$

→ **evolving (increasing) blazar population consistent with observed declining evolution (*Fermi* flux limit)!**



# How many TeV blazars are there at high- $z$ ?




Hopkins+ (2007)



# Fermi probes “dragons” of the gamma-ray sky

## Fermi LAT Extragalactic Gamma-ray Background



# Extragalactic gamma-ray background

- assume all TeV blazars have identical intrinsic spectra:

$$F_E = L\hat{F}_E \propto \frac{1}{(E/E_b)^{\alpha_L-1} + (E/E_b)^{\alpha-1}},$$

$E_b$  is break energy,

$\alpha_L < \alpha$  are low and high-energy spectral indexes

- extragalactic gamma-ray background (EGRB):

$$E^2 \frac{dN}{dE}(E, z) = \frac{1}{4\pi} \int_z^\infty dV(z') \frac{\eta_B \tilde{\Lambda}_Q(z') \hat{F}_{E'}}{4\pi D_L^2} e^{-\tau_E(E', z')},$$

$E' = E(1 + z')$  is gamma-ray energy at *emission*,


$\tilde{\Lambda}_Q$  is physical quasar luminosity density,

$\eta_B \sim 0.2\%$  is blazar fraction,  $\tau$  is optical depth





# Extragalactic gamma-ray background



Broderick, Chang, C.P. (2012)

- *dotted*: unabsorbed EGRB due to TeV blazars
- *dashed*: absorbed EGRB due to TeV blazars
- *solid*: absorbed EGRB, after subtracting the resolved TeV blazars ( $z < 0.25$ )




# Outline

- 1 Physics of blazar heating
  - TeV emission from blazars
  - Plasma instabilities and magnetic fields
  - Extragalactic gamma-ray background
- 2 The intergalactic medium
  - Properties of blazar heating
  - Thermal history of the IGM
  - The Lyman- $\alpha$  forest
- 3 Structure formation
  - Formation of dwarf galaxies
  - Puzzles in galaxy formation
  - Conclusions



# Evolution of the heating rates



Chang, Broderick, C.P. (2012)

# Blazar heating vs. photoheating

- total power from AGN/stars vastly exceeds the TeV power of blazars



# Blazar heating vs. photoheating

- total power from AGN/stars vastly exceeds the TeV power of blazars
- $T_{\text{IGM}} \sim 10^4$  K (1 eV) at mean density ( $z \sim 2$ )

$$\varepsilon_{\text{th}} = \frac{kT}{m_p c^2} \sim 10^{-9}$$



# Blazar heating vs. photoheating

- total power from AGN/stars vastly exceeds the TeV power of blazars
- $T_{\text{IGM}} \sim 10^4$  K (1 eV) at mean density ( $z \sim 2$ )

$$\varepsilon_{\text{th}} = \frac{kT}{m_p c^2} \sim 10^{-9}$$

- radiative energy ratio emitted by BHs in the Universe (Fukugita & Peebles 2004)

$$\varepsilon_{\text{rad}} = \eta \Omega_{\text{bh}} \sim 0.1 \times 10^{-4} \sim 10^{-5}$$



# Blazar heating vs. photoheating

- total power from AGN/stars vastly exceeds the TeV power of blazars
- $T_{\text{IGM}} \sim 10^4$  K (1 eV) at mean density ( $z \sim 2$ )

$$\varepsilon_{\text{th}} = \frac{kT}{m_p c^2} \sim 10^{-9}$$

- radiative energy ratio emitted by BHs in the Universe (Fukugita & Peebles 2004)

$$\varepsilon_{\text{rad}} = \eta \Omega_{\text{bh}} \sim 0.1 \times 10^{-4} \sim 10^{-5}$$

- fraction of the energy energetic enough to ionize H I is  $\sim 0.1$ :

$$\varepsilon_{\text{UV}} \sim 0.1 \varepsilon_{\text{rad}} \sim 10^{-6} \quad \rightarrow \quad kT \sim \text{keV}$$



# Blazar heating vs. photoheating

- total power from AGN/stars vastly exceeds the TeV power of blazars
- $T_{\text{IGM}} \sim 10^4$  K (1 eV) at mean density ( $z \sim 2$ )

$$\varepsilon_{\text{th}} = \frac{kT}{m_p c^2} \sim 10^{-9}$$

- radiative energy ratio emitted by BHs in the Universe (Fukugita & Peebles 2004)

$$\varepsilon_{\text{rad}} = \eta \Omega_{\text{bh}} \sim 0.1 \times 10^{-4} \sim 10^{-5}$$

- fraction of the energy energetic enough to ionize H I is  $\sim 0.1$ :

$$\varepsilon_{\text{UV}} \sim 0.1 \varepsilon_{\text{rad}} \sim 10^{-6} \quad \rightarrow \quad kT \sim \text{keV}$$

- photoheating efficiency  $\eta_{\text{ph}} \sim 10^{-3} \quad \rightarrow \quad kT \sim \eta_{\text{ph}} \varepsilon_{\text{UV}} m_p c^2 \sim \text{eV}$   
(limited by the abundance of H I/He II due to the small recombination rate)





# Blazar heating vs. photoheating

- total power from AGN/stars vastly exceeds the TeV power of blazars
- $T_{\text{IGM}} \sim 10^4$  K (1 eV) at mean density ( $z \sim 2$ )

$$\varepsilon_{\text{th}} = \frac{kT}{m_p c^2} \sim 10^{-9}$$

- radiative energy ratio emitted by BHs in the Universe (Fukugita & Peebles 2004)

$$\varepsilon_{\text{rad}} = \eta \Omega_{\text{bh}} \sim 0.1 \times 10^{-4} \sim 10^{-5}$$

- fraction of the energy energetic enough to ionize H I is  $\sim 0.1$ :

$$\varepsilon_{\text{UV}} \sim 0.1 \varepsilon_{\text{rad}} \sim 10^{-6} \rightarrow kT \sim \text{keV}$$

- photoheating efficiency  $\eta_{\text{ph}} \sim 10^{-3} \rightarrow kT \sim \eta_{\text{ph}} \varepsilon_{\text{UV}} m_p c^2 \sim \text{eV}$


(limited by the abundance of H I/He II due to the small recombination rate)

- blazar heating efficiency  $\eta_{\text{bh}} \sim 10^{-3} \rightarrow kT \sim \eta_{\text{bh}} \varepsilon_{\text{rad}} m_p c^2 \sim 10 \text{ eV}$

(limited by the total power of TeV sources)



# Thermal history of the IGM




Chang, Broderick, C.P. (2012)




# Evolution of the temperature-density relation

no blazar heating



# Evolution of the temperature-density relation


no blazar heating



- blazars and extragalactic background light are uniform:  
→ blazar heating rate independent of density

# Evolution of the temperature-density relation

no blazar heating




- blazars and extragalactic background light are uniform:
  - blazar heating rate independent of density
  - makes low density regions *hot*
  - causes inverted temperature-density relation,  $T \propto 1/\delta$




# Evolution of the temperature-density relation

no blazar heating



with blazar heating




Chang, Broderick, C.P. (2012)

- blazars and extragalactic background light are uniform:
  - blazar heating rate independent of density
  - makes low density regions *hot*
  - causes inverted temperature-density relation,  $T \propto 1/\delta$




# Blazars cause hot voids

no blazar heating




with blazar heating




Chang, Broderick, C.P. (2012)

# Blazars cause hot voids

no blazar heating



with blazar heating



Chang, Broderick, C.P. (2012)

- blazars completely change the thermal history of the diffuse IGM and late-time structure formation




# Simulations with blazar heating

Puchwein, C.P., Springel, Broderick, Chang (2012):

- $L = 15h^{-1}\text{Mpc}$  boxes with  $2 \times 384^3$  particles
- one reference run without blazar heating
- three with blazar heating at different levels of efficiency  
(address uncertainty)
- used an up-to-date model of the UV background (Faucher-Giguère+ 2009)




# Temperature-density relation



Puchwein, C.P., Springel, Broderick, Chang (2012)




# Ly- $\alpha$ spectra





Puchwein+ (2012)



# Optical depths and temperatures



# Optical depths and temperatures




Puchwein+ (2012)


Redshift evolutions of effective optical depth and IGM temperature match data only with additional heating, e.g., provided by blazars!




# Ly- $\alpha$ flux PDFs and power spectra



Puchwein+ (2012)




# Voigt profile decomposition



- decomposing Lyman- $\alpha$  forest into individual Voigt profiles
- allows studying the thermal broadening of absorption lines

# Voigt profile decomposition – line width distribution



Puchwein+ (2012)



# Lyman- $\alpha$ forest in a blazar heated Universe

improvement in modelling the Lyman- $\alpha$  forest is a direct consequence of the peculiar properties of blazar heating:

- **heating rate independent of IGM density**  $\rightarrow$  naturally produces the inverted  $T-\rho$  relation that Lyman- $\alpha$  forest data demand
- **recent and continuous nature of the heating** needed to match the redshift evolutions of all Lyman- $\alpha$  forest statistics
- **magnitude of the heating rate required by Lyman- $\alpha$  forest data**  
 **$\sim$  the total energy output of TeV blazars** (or equivalently  $\sim 0.2\%$  of that of quasars)




# Outline

- 1 Physics of blazar heating
  - TeV emission from blazars
  - Plasma instabilities and magnetic fields
  - Extragalactic gamma-ray background
- 2 The intergalactic medium
  - Properties of blazar heating
  - Thermal history of the IGM
  - The Lyman- $\alpha$  forest
- 3 **Structure formation**
  - **Formation of dwarf galaxies**
  - **Puzzles in galaxy formation**
  - **Conclusions**




# Entropy evolution

## temperature evolution




# Entropy evolution

temperature evolution



entropy evolution



C.P., Chang, Broderick (2012)

- evolution of entropy,  $K_e = kTn_e^{-2/3}$ , governs structure formation
- blazar heating: late-time, evolving, modest entropy floor



# Dwarf galaxy formation – Jeans mass

- thermal pressure opposes gravitational collapse on small scales
- characteristic length/mass scale below which objects do not form



# Dwarf galaxy formation – Jeans mass

- thermal pressure opposes gravitational collapse on small scales
- characteristic length/mass scale below which objects do not form
- hotter IGM  $\rightarrow$  higher IGM pressure  $\rightarrow$  **higher Jeans mass:**

$$M_J \propto \frac{c_s^3}{\rho^{1/2}} \propto \left( \frac{T_{\text{IGM}}^3}{\rho} \right)^{1/2} \rightarrow \frac{M_{J,\text{blazar}}}{M_{J,\text{photo}}} \approx \left( \frac{T_{\text{blazar}}}{T_{\text{photo}}} \right)^{3/2} \gtrsim 30$$

$\rightarrow$  depends on instantaneous value of  $c_s$



# Dwarf galaxy formation – Jeans mass

- thermal pressure opposes gravitational collapse on small scales
- characteristic length/mass scale below which objects do not form
- hotter IGM  $\rightarrow$  higher IGM pressure  $\rightarrow$  **higher Jeans mass:**


$$M_J \propto \frac{c_s^3}{\rho^{1/2}} \propto \left( \frac{T_{\text{IGM}}^3}{\rho} \right)^{1/2} \rightarrow \frac{M_{J,\text{blazar}}}{M_{J,\text{photo}}} \approx \left( \frac{T_{\text{blazar}}}{T_{\text{photo}}} \right)^{3/2} \gtrsim 30$$

$\rightarrow$  depends on instantaneous value of  $c_s$

- “**filtering mass**” depends on full thermal history of the gas: accounts for delayed response of pressure in counteracting gravitational collapse in the expanding universe
- apply corrections for **non-linear collapse**



# Dwarf galaxy formation – Filtering mass




C.P., Chang, Broderick (2012)






# Peebles' void phenomenon explained?

mean density



void,  $1 + \delta = 0.5$




C.P., Chang, Broderick (2012)

- blazar heating efficiently suppresses the formation of void dwarfs within existing DM halos of masses  $< 3 \times 10^{11} M_\odot$  ( $z=0$ )
- may reconcile the number of void dwarfs in simulations and the paucity of those in observations




# “Missing satellite” problem in the Milky Way




Substructures in cold DM simulations much more numerous than observed number of Milky Way satellites!

# When do dwarfs form?




Dolphin+ (2005)



isochrone fitting for different metallicities  $\rightarrow$  star formation histories



# When do dwarfs form?




Dolphin+ (2005)




# Milky Way satellites: formation history and abundance

## satellite formation time




# Milky Way satellites: formation history and abundance

satellite formation time



satellite luminosity function





Maccio+ (2010)

- blazar heating suppresses late satellite formation, may reconcile low observed dwarf abundances with CDM simulations



# Galactic H I-mass function

Mo+ (2005)



- H I-mass function is too flat (i.e., gas version of missing dwarf problem!)
- photoheating and SN feedback too inefficient
- IGM entropy floor of  $K \sim 15 \text{ keV cm}^2$  at  $z \sim 2 - 3$  successful!



# Conclusions on blazar heating

- explains puzzles in high-energy astrophysics:
  - lack of GeV bumps in blazar spectra without IGM  $B$ -fields
  - *unified TeV blazar-quasar model* explains Fermi source counts and extragalactic gamma-ray background





# Conclusions on blazar heating

- explains puzzles in high-energy astrophysics:
  - lack of GeV bumps in blazar spectra without IGM  $B$ -fields
  - *unified TeV blazar-quasar model* explains Fermi source counts and extragalactic gamma-ray background
- novel mechanism; dramatically alters thermal history of the IGM:
  - uniform and  $z$ -dependent preheating
  - rate independent of density  $\rightarrow$  inverted  $T-\rho$  relation
  - quantitative self-consistent picture of high- $z$  Lyman- $\alpha$  forest




# Conclusions on blazar heating

- explains puzzles in high-energy astrophysics:
  - lack of GeV bumps in blazar spectra without IGM  $B$ -fields
  - *unified TeV blazar-quasar model* explains Fermi source counts and extragalactic gamma-ray background
- novel mechanism; dramatically alters thermal history of the IGM:
  - uniform and  $z$ -dependent preheating
  - rate independent of density  $\rightarrow$  inverted  $T-\rho$  relation
  - quantitative self-consistent picture of high- $z$  Lyman- $\alpha$  forest
- significantly modifies late-time structure formation:
  - suppresses late dwarf formation (in accordance with SFHs): “missing satellites”, void phenomenon, H I-mass function
  - group/cluster bimodality of core entropy values




# When do clusters form?

mass accretion history



mass accretion rates




C.P., Chang, Broderick (2012)

- most cluster gas accretes after  $z = 1$ , when blazar heating can have a large effect (for late forming objects)!



# Entropy floor in clusters


## Cluster entropy profiles



Cavagnolo+ (2009)


# Entropy floor in clusters

## Cluster entropy profiles



Cavagnolo+ (2009)

## Planck stacking of optical clusters




Planck Collaboration (2011)

- Do optical and X-ray/Sunyaev-Zel'dovich cluster observations probe the same population? (Hicks+ 2008, Planck Collaboration 2011)




# Entropy profiles: effect of blazar heating

varying formation time



varying cluster mass




C.P., Chang, Broderick (2012)

**assume** big fraction of intra-cluster medium collapses from IGM:

- redshift-dependent entropy excess in cores
- greatest effect for late forming groups/small clusters



# Gravitational reprocessing of entropy floors




Borgani+ (2005)

- greater initial entropy  $K_0$   
→ more shock heating  
→ greater increase in  $K_0$   
over entropy floor
- net  $K_0$  amplification of 3-5
- expect:  
median  $K_{e,0} \sim 150 \text{ keV cm}^2$   
max.  $K_{e,0} \sim 600 \text{ keV cm}^2$




# Cool-core versus non-cool core clusters






# Cool-core versus non-cool core clusters




# Cool-core versus non-cool core clusters




- time-dependent preheating + gravitational reprocessing  
→ CC-NCC bifurcation (two attractor solutions)
- need hydrodynamic simulations to confirm this scenario


# How efficient is heating by AGN feedback?




# How efficient is heating by AGN feedback?




# How efficient is heating by AGN feedback?




# How efficient is heating by AGN feedback?



# How efficient is heating by AGN feedback?




# How efficient is heating by AGN feedback?






# How efficient is heating by AGN feedback?



# How efficient is heating by AGN feedback?



**AGNs cannot transform CC to NCC clusters** (on a buoyancy timescale)

